

# **INDIAN ASSOCIATION OF PHYSICS TEACHERS**

# NATIONAL STANDARD EXAMINATION IN CHEMISTRY (NSEC) 2019-20

Examination Date : 24-11-2019

Time: 2 Hrs.

Max. Marks : 240

# Q. PAPER CODE: 34

HBCSE Olympiad (STAGE - 1)

Write the question paper code mentioned above on YOUR answer sheet (in the space provided), otherwise your answer sheet will NOT be assessed. Note that the same Question paper. Code appears on each page of the question paper.

## INSTRUCTION TO CANDIDATES

- 1. Use of mobile phones, smart watches and ipads during examination is **STRICTLY PROHIBITED.**
- 2. In addition to this question paper, you are given answer sheet along with Candidate's copy.
- 3. On the answer sheet, make all the entries carefully in the space provided ONLY in **BLOCK** CAPITALS as well as by properly darkening the appropriate bubbles.
- Incomplete/Incorrect/carelessly filled information may disqualify your candidature.
- 4. On the answer sheet, use only **BLUE or BLACK BALL POINT PEN** for making entries and filling the bubbles.
- 5. Your **ten-digit roll number and date of birth** entered on the OMR Answer Sheet shall remain your login credentials means login id and password respectively for accessing your performance / result in NSE-2019.
- 6. Question paper contain 80 multiple-choice question. Each question has 4 options, out of which only one is correct. Choose the correct alternative and fill the appropriate bubble, as shown



- 7. A correct answer carries 3 marks and 1 mark will be deducted for each wrong answer.
- 8. Any rough work should be done only in the space provided.
- 9. Use of **non-programmable scientific** calculator is allowed.
- 10. No candidate should leave the examination hall before the completion of the examination.
- 11. After submitting your answer paper, take away the Candidate's copy for your reference.

Please DO NOT make any mar other than filling the appropriate bubbles properly in the space provided on the answer sheet. Answer sheet are evaluated using machine, hence CHANGE OF ENTRY IS NOT ALLOWED.

Scratching or overwriting may result in wrong score.

DO NOT WRITE ANYTHING ON THE BACK OF ANSWER SHEET.

### Instructions to candidates :

- You may read the following instructions after submitting the answer sheet.
- 12. Comments/Inquiries/Grievences regarding this quesiton paper, if any, can be shared on the Inquiry/Grievence column on <u>www.iaptexam.in</u> on the specified format till 27<sup>th</sup> November, 2019.
- 13. The answers/solutions to this question paper will be available on the website : www.iapt.org.in by 2<sup>nd</sup> December, 2019.

## 14. CERTIFICATES and AWARDS –

- Following certificates are awarded by IAPT/ACT to students successful in the NSEC-2019. (i) "CENTRE TOP 10%"
  - (ii) "STATE TOP 1%"
    - (iii) "NATIONAL TOP 1%"
  - (iv) "GOLD MEDAL & MERIT CERTIFICATE" to all students who attend OCSC-2020 at HBCSE.
- 15. All these certificates (except GOLD Medal) will be sent/dispatched to the centre in-charge by February 1, 2020 along with the result sheet of the centre.
- 16. List of students (with centre number and roll number only) having score above MAS will be displayed on the website : www.iapt.org.in by December 20<sup>nd</sup>, 2019. See the Minimum Admissible score Clause on the Student's brochure on the web.
- 17. List of Students eligible for the Indian National Chemistry Olympiad (INChO-2020) shall be displayed on <u>www.iapt.org.in</u> by December 28,2019. These students have to register/enroll themselves on the website : Olympiads.hbcse.tifr.in of HBCSE Mumbai within the specified period.

# **Resonance Eduventures Limited**

REGISTERED & CORPORATE OFFICE : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph.No.: 0744-2777777, 0744-2777700 | Toll Free: 1800 258 5555 | FAX No.: +91-022-39167222 | To Know more: sms RESO at 56677

 Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

 Toll Free : 1800 258 5555
 73400 10333
 If facebook.com/ResonanceEdu
 Ituiter.com/ResonanceEdu
 Ituiter.com/ResonanceEdu
 Ituiter.com/ResonanceEdu

This solution was download from Resonance Solution portal



Total Time : 120 minutes (A–1 and A–2)



2. The following Ellingham diagram depicts the oxidation of 'C', 'CO' and Fe'. Which of the following is correct?



FeO can be reduced by CO below 600 K Π. FeO can be reduced by C above 1000 K III. FeO can be reduced by CO above 1000 K IV. (A) II and III (B) I and IV (C) I and III (D) II and IV Ans. (A) Sol. Below 600 K, reaction- $FeO + CO \longrightarrow Fe + CO_2$  is feasible Above 1000 K, reaction- $FeO + C \longrightarrow Fe + CO$  is feasible 3. A balance having a precision of 0.0001 g was used to measure a mass of a sample of about 15 g. The number of significant figures to be reported in this measurement is (A) 2 (B) 3 (C) 5 (D) 1 Ans. (C)

Kesonance         Website : www.resonance.ac.in   E-mail : contact@resonance.ac.in         NSEC24112019 # 2           Toll Free : 1800 258 5555   CIN: U80302RJ2007PLC024029         NSEC24112019 # 2		A Resonance®	Reg. & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar R	oad, Kota (Raj.)- 324005
Educating for better tomorrow         Toll Free : 1800 258 5555   CIN: U80302RJ2007PLC024029         RSEC24112019 # 2			Website : www.resonance.ac.in   E-mail : contact@resonance.ac.in	NSEC24112010 # 2
	Educating for better tomorrow	Toll Free : 1800 258 5555   CIN: U80302RJ2007PLC024029	N3EC24112019#2	



Resonance<sup>®</sup> NATIONAL STANDARD EXAMINATION IN CHEMISTRY (NSEC) 2019-20 | 24-11-2019 7. An electrochemical cell was constructed with Fe<sup>2+</sup>/Fe and Cd<sup>2+</sup>/Cd at 25°C with initial concentrations of  $[Fe^{2+}] = 0.800$  M and  $[Cd^{2+}] = 0.250$  M. The EMF of the cell when  $[Cd^{2+}]$  becomes 0.100 M is Half cell E°(V) Fe<sup>2+</sup>(aq)/Fe(s) -0.44-0.40Cd<sup>2+</sup>(aq)/Cd(s) (A) 0.013 V (B) 0.011 V (C) 0. 051 V (D) 0.022 V Ans. (B) Sol. Reaction  $Cd^{2+} + Fe(s) \longrightarrow Fe^{2+} + Cd(s)$ 0.25 M t = 00.8 M t = t0.25 - x0.8 + x= 0.1∴ x = 0.15. At this instance  $[Cd^{2+}] = 0.1 M$  $[Fe^{2+}] = 0.8 + x = 0.95$  $E_{cell} = E_{cell}^{0} - \frac{0.059}{2} \log \frac{[Fe^{2+}]}{[Cd^{2+}]}$  $E_{cell} = (-0.4 + 0.44) - \frac{0.059}{2} \log \left(\frac{0.95}{0.1}\right)$ = 0.011V 8. The kinetic energy of the photoelectrons ejected by a metal surface increased from 0.6 eV to 0.9 eV when the energy of the incident photons was increased by 20%. The work function of the metal is (A) 0.66 eV (B) 0.72 eV (C) 0.90 eV (D) 0.30 eV Ans. (C) Sol.  $KE_{max} = E - E_0$  $0.6 = E - E_0 \implies$  $E = 0.6 + E_0$  $0.9 = 1.2E - E_0 \implies 1.2E = 0.9 + E_0$ On dividing  $\frac{1}{1.2} = \frac{0.6 + E_0}{0.9 + E_0}$  $0.9 + E_0 = 0.72 + 1.2 E_0$  $\therefore E_0 = 0.9 \text{ eV}$ 9. The alkene ligand  $(\pi - C_2R_4)$  is both a ' $\sigma$ ' donoar and a ' $\pi$ ' acceptor, similar to the CO ligand in metal carbonyls, and exhibits synergic bonding with metals. Correct order of C-C bond length in K[PtCl<sub>3</sub>( $\pi$ –C<sub>2</sub>R<sub>4</sub>)] complexes in which R = H, F or CN is (A) H > F > CN (B) H > CN > F(C) CN > F > H(D) F > H > CN(C) Ans.  $K[PtCl_3(\pi - C_2R_4]]$ Sol. In C<sub>2</sub>R<sub>4</sub> as electron withdrawing nature of R<sup> $\uparrow$ </sup>  $\Rightarrow$  back bonding from pt to alkene<sup> $\uparrow$ </sup>  $\Rightarrow$  C–C bond order $\downarrow$ so order of electron withdrawing nature-H < F < CNorder of C-C bond length  $K[PtCl_3(\pi - C_2R_4)] < K[PtCl_3(\pi - C_2F_4)] < K[PtCl_3(\pi - C_2(CN)_4)]$ The correct order of CFSE among [Zn(NH<sub>3</sub>)<sub>4</sub>]<sup>2+</sup> and [Co(NH<sub>3</sub>)<sub>6</sub>]<sup>2+</sup> and [Co(NH<sub>3</sub>)<sub>6</sub>]<sup>3+</sup> is 10. (B)  $[Zn(NH_3)_4]^{2+} > [Co(NH_3)_6]^{2+} > [Co(NH_3)_6]^{3+}$ (D)  $[Co(NH_3)_6]^{2+} > [Co(NH_3)_6]^{3+} > [Zn(NH_3)_4]^{2+}$ (A)  $[Co(NH_3)_6]^{3+} > [Co(NH_3)_6]^{2+} > [Zn(NH_3)_4]^{2+}$ (C)  $[Co(NH_3)_6]^{3+} > [Zn(NH_3)_4]^{2+} > [Co(NH_3)_6]^{2+}$ Ans. (A) CFSE in [Zn(NH<sub>3</sub>)<sub>4</sub>]<sup>2+</sup> is zero. Sol. so correct order of CFSE  $Co(NH_3)_{6}^{3+} > Co(NH_3)_{6}^{2+} > [Zn(NH_3)_4]^{2+}$ 



 Reg. & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.)- 324005

 Website: www.resonance.ac.in | E-mail : contact@resonance.ac.in

 Toll Free : 1800 258 5555 | CIN: U80302RJ2007PLC024029

NSEC24112019 # 4







	Reg. & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar R	oad, Kota (Raj.)- 324005
Resonance	Website : www.resonance.ac.in   E-mail : contact@resonance.ac.in	NSEC24112010 # 7
Educating for better tomorrow	Toll Free : 1800 258 5555   CIN: U80302RJ2007PLC024029	N3EC24112019#1

## Resonance<sup>®</sup> I NATIONAL STANDARD EXAMINATION IN CHEMISTRY (NSEC) 2019-20 | 24-11-2019

- 21. The set in which all the species are diamagnetic is
  - (A) B<sub>2</sub>, O<sub>2</sub>, NO (B) O<sub>2</sub>, O<sub>2</sub><sup>+</sup>, CO (D) C<sub>2</sub>, O<sub>2</sub><sup>2-</sup>, NO<sup>+</sup> (C) N<sub>2</sub>, O<sub>2</sub><sup>-</sup>, CN<sup>-</sup>
- Ans. (D)
- Sol. B<sub>2</sub>, O<sub>2</sub>, NO, O<sub>2</sub><sup>+</sup>, O<sub>2</sub><sup>-</sup>  $\rightarrow$  Paramagnetic
  - $O_2^{2-}$ ,  $C_2$ ,  $N_2$ ,  $NO^+$ ,  $CO \rightarrow$  Diamagnetic
- 22. A solid comprises of three types of elements, 'P', 'Q' and 'R'. 'P' forms an FCC lattice in which 'Q' and 'R' occupy all the tetrahedral voids and half the octahedral voids respectively. The molecular formula of the solid is
  - (A)  $P_2Q_2R$ (B)  $PQ_2R_4$
  - (C)  $P_4Q_2R$ (D) P<sub>4</sub>QR
- Ans. (A)
- Sol. Formula of solid

 $= P_4 Q_8 R_{4/2} = P_4 Q_8 R_2 = P_2 Q_4 R$ 

23. The following qualitative plots depict the first, second and third ionization energies (I.E.) of Mg, Al and K. Among the following, the correct match of I.E. and the metal is



## Resonance<sup>®</sup> Examination IN CHEMISTRY (NSEC) 2019-20 | 24-11-2019

#### 24. The structure of compound 'X' (C<sub>8</sub>H<sub>11</sub>NO) based on the following tests and observations is

Reagent/s	Observation
Neutral FeCl <sub>3</sub>	No coloration
Lucas reagent	Turbidity
NaNO <sub>2</sub> /HCl at 273 K	Yellow oil



Toll Free : 1800 258 5555 | CIN: U80302RJ2007PLC024029

Educating for better tomorrow

NSEC24112019 # 9

Educ		DARD EXAMINATION IN CHEMISTRY (NSEC) 2019-20   24-11-2019				
26.	Reaction of $C_6H_5MgBr$ with phenol g	gives				
	(A)	(В) ОН				
		(D) Br				
Ans.	(A)					
Sol.	PhMgBr + HO	) + PhOMgBr				
27.	The power and wavelength emitted are 1.0 mW and 670 nm respect presentation of 5 minutes is (A) $1.01 \times 10^9$ (C) $1.6 \times 10^{16}$	by a laser pointer commonly used in Power Point presentations tively. Number of photons emitted by this pointer during a (B) $1.01 \times 10^{21}$ (D) $1.01 \times 10^{18}$				
Ans.	(D)					
Sol.	Total energy emitted in 5 min = $10^{-3}$	$3 \times 5 \times 60 \text{ J} = 0.3 \text{ J}$				
	$E = \frac{Nhc}{\lambda}$					
	$N = \xrightarrow{E \times \lambda}_{hc} = \xrightarrow{0.3 \times 670 \times 10^{-9}}_{6.626 \times 10^{-34} \times 3 \times 10^8} = 1.$	01 × 10 <sup>18</sup>				
28.	The work done (kJ) in the irreversibl bar to 100 bar at 25°C at constant e (A) 2452	e isothermal compression of 2.0 moles of an ideal gas from 1 xternal pressure of 500 bar is (B) 490				
	(C) 2486	(D) – 490				
Ans.	(A)					
Sol.	$W = - P_{ext} (\Delta V)$					
	$= -500 \left(\frac{nRT}{100} - \frac{nRT}{1}\right)$					
	= – nRT (5 – 500)					
	= + 2 × 8.314 × 298 × 495 × 10 <sup>-3</sup> kJ	= + 2452 kJ				
29.	Atropine ( $C_{17}H_{23}O_3N$ ) is a naturally of The degree of unsaturation in atropi (A) 7	occurring compound used to treat certain types of poisoning. ne is (B) 6				
	(C) 5	(D) 4				
Ans.	(A)					
	Reg. & Corp. C	office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.)- 324005				
$\left  \right\rangle $	Contract @resonance.ac.in       E-mail : contact @resonance.ac.in       NSEC24112019 # 10         Toll Free : 1800 258 5555   CIN: U80302RJ2007PLC024029       NSEC24112019 # 10					

**Sol.** (C<sub>17</sub>H<sub>23</sub>O<sub>3</sub>N)  

$$DU = \frac{2N+2+Z-M-X}{2}$$
Where = N = no. of carbon  
M = no. of H  
Z = no. of N  
X = no. of Halogen  
Hence = DU = 7  
**30.** MnCl<sub>2</sub>.4H<sub>2</sub>O (molar mass = 198 g mol<sup>-1</sup>) when dissolved in water forms a complex of Mn<sup>2+</sup>. An  
aquecus solution containing 0.400 g of MnCl<sub>2</sub>.4H<sub>2</sub>O was passed through a column of a cation  
exchange resin and the acid solution coming out was neutralized with 10 mL of 0.20 M NaOH. The  
formula of the complex formed is  
(A) [Mn(H<sub>2</sub>O)<sub>2</sub>Cl<sub>2</sub>] (B) [Mn(H<sub>2</sub>O)<sub>3</sub>Cl<sub>3</sub>]  
**Ans.** (C)  
**Sol.** MCl<sub>2</sub>.4H<sub>2</sub>O  $\longrightarrow$  xH<sup>+</sup>  $\xrightarrow{NOH}$   
moles of Compound taken =  $\frac{0.4}{198}$   
moles of OH<sup>+</sup> required = 0.2 × 0.01 = 0.002  
 $\therefore \frac{x \times 0.4}{198} = 0.002$   
 $\Rightarrow x = 1$   
Compound is [Mn(H<sub>2</sub>O)<sub>2</sub>Cl]Cl  
**31.** Which of the following is NOT correct about hydrides ?  
I. Saline hydrides are stoichiometric and metallic hydrides re non-stoichiometric  
II. BeH<sub>2</sub> is monomeric whereas MgH<sub>2</sub> is polymeric  
III. Hydrides of the elements of Group 13 are electron deficient and those of Group 15 are electron  
rich  
V. NaH reacts with water and liberates H<sub>2</sub> whereas B<sub>3</sub>H<sub>6</sub> does not react with water  
(A) IV only (B) I and III

(D) II and IV (C) III only

Ans. (D)



**Sol.** BeH<sub>2</sub> is polymetric while MgH<sub>2</sub> is monomeric

 $2NaH + H_2O \longrightarrow 2NaOH + H_2$ 

 $B_2H_6 + H_2O \longrightarrow B(OH)_3 + H_2$ 

32. The compounds 'X' and 'Y' formed in the following reaction are



(A) hemiacetals with identical physical and chemical properties

- (B) acetals with identical physical and chemical properties
- (C) hemiacetals with different physical and chemical properties
- (D) acetals with different physical and chemical properties

Ans. (C)



x and y are hemiacetal and are diastereisomers of each other, so they will have different physical and chemical properties.

- **33.** Aqueous solution of slaked lime, Ca(OH)<sub>2</sub>, is extensively used in municipal waste water treatment. Maximum pH possible in an aqueous solution of slaked lime is  $(K_{sp} \text{ of } Ca(OH)_2 = 5.5 \times 10^{-6})$ (A) 1.66 (B) 8.14
  - (C) 12.04

(D) 12.34

Ans. (D)

Sol.

S 2S

 $Ca(OH)_2(s) \Longrightarrow Ca^{2+} + 2OH^-$ 

 $K_{sp} = 4S^3$ ; S = 0.011

∴ [OH<sup>-</sup>] = 2S = 0.022

 $P_{OH} = 1.65$ ; =  $P_{H} = 12.34$ 



	Reg. & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar R	load, Kota (Raj.)- 324005	
Kesonance	Website : www.resonance.ac.in   E-mail : contact@resonance.ac.in	NSEC24112010 # 12	
Educating for better tomorrow	Toll Free : 1800 258 5555   CIN: U80302RJ2007PLC024029	NOLC24112019#12	

## 

34.	An electron present in the third exited state of a H atom returns of the first exited state and then to the ground state. If $\lambda_1$ and $\lambda_2$ are the wavelengths of light emitted in these two transitions					
	respectively, $\lambda_1 : \lambda_2$ is					
	(A) 4 : 1		(B) 5 : 9			
<b>A</b>	(C) 3 : 1		(D) 2 : 1			
Ans.	(A)					
0	n = 4					
501.	$\frac{\Rightarrow \lambda_1}{n=3}$					
	n = 2					
	$\rightarrow \lambda_2$ n = 1					
	$\frac{1}{\lambda_1} = R_{H} \left( \frac{1}{4} - \frac{1}{16} \right) \Longrightarrow \lambda_1 = \frac{16}{3R_{H}}$					
	1 (1 1) 4					
	$\frac{1}{\lambda_2} = R_H \left( \frac{1}{1} - \frac{1}{4} \right) \Rightarrow \lambda_2 = \frac{1}{3R_H}$					
	λ <sub>1</sub> 16 4					
	$\therefore \frac{1}{\lambda_2} = \frac{1}{4} = \frac{1}{1}$					
35	The perceptage dissociation of	0.08 M aqueous	s acetic acid solution at 25ºC is (	K <sub>2</sub> of acetic acid at		
	$25^{\circ}C = 1.8 \times 10^{-5}$ )	0.00 11 494004				
	(A) 2.92		(B) 1.5			
Anc	(C) 1.2 (B)		(D) 4.8			
Sol.	$CH_3COOH \longrightarrow CH_3COO^- + F$	l⊕				
	C					
	$C(1-\alpha)$ $C\alpha$ $C\alpha$	ι				
	$\alpha = \sqrt{\frac{K_a}{K_a}} = \sqrt{\frac{1.8 \times 10^{-5}}{1.8 \times 10^{-5}}} = 0.015$					
	VC V 0.08					
	% dissociation = 1.5					
36.	In which of the following, is a ne	w C-C bond for	med in the product ?			
	I. CH <sub>3</sub> CHO $\xrightarrow{\text{dil. NaOH}}$					
	II. CH <sub>3</sub> MgCl + C <sub>2</sub> H <sub>5</sub> OH <u>heat</u>	→				
	III. CO <sub>2</sub> + CH <sub>3</sub> MgBr $\xrightarrow{H_3O^+}$					
	IV. C <sub>2</sub> H <sub>2</sub> + NaNH <sub>2</sub> $\xrightarrow{CH_3Br}$					
	(A) I, III and IV (B) II a	nd III	(C) III only	(D) III and IV		
Ans.	(A)	04 0				
Sol.	(I) $CH_3CH = O \xrightarrow{dil/NaOH}_{Aldol} CH_3$	_СH–СH₂–С–Н (С	–C bond formed)			
	(II) CH <sub>3</sub> MgBr + EtOH $\xrightarrow{\text{heat}}_{\text{acid, base}}$	(CH <sub>4</sub> + EtOMgB	r (No C–C bond)			
		UH3UUUH (U-U	, bona)			
	(IV) $CH \equiv CH + NaNH_2 \longrightarrow d$	сцё CH <sub>3</sub> -Br	(C–C bond)			
	. ,					
0	Reg. & C	orp. Office: CG Towe	r, A-46 & 52, IPIA, Near City Mall, Jhalawar F	≀oad, Kota (Raj.)- 324005		
$\square$	Educating for better tomorrow Website	: www.resonance.ac.ir : 1800 258 5555   <b>CIN:</b> U	<b>E-mail :</b> contact@resonance.ac.in 80302RJ2007PLC024029	NSEC24112019 # 13		











	Atting for better temorrow NATIONAL STANDARD EXAMINATION IN CHEMISTRY (NSEC) 2019-20   24-11-2019				
55.	Mercury is highly hazardous and hence its concentration is expressed in the units of ppb (micrograms of Hg present in 1 L of water). Permissible level of Hg in drinking water is 0.0335 ppb. Which of the following is an alternate representation of this concentration? (A) $3.35 \times 10^{-2}$ mg dm <sup>-3</sup> (B) $3.35 \times 10^{-5}$ mg dm <sup>-3</sup> (C) $3.35 \times 10^{-5}$ mg m <sup>-3</sup> (D) $3.35 \times 10^{-4}$ g L <sup>-1</sup>				
Ans. Sol.	(B) Concentration of Hg = 0.0335 ppb of Hg( $\mu$ g/L) = 0.0335 × 10 <sup>-3</sup> mg/lt = 3.35 × 10 <sup>-5</sup> mg/dm <sup>3</sup>				
56.	The correct sequence of reaction which will yield 4-nitrobenzoic acid from benzene is(A) CH <sub>3</sub> Cl; HNO <sub>3</sub> /H <sub>2</sub> SO <sub>4</sub> ; KMnO <sub>4</sub> /OH <sup>-</sup> (B) HNO <sub>3</sub> /H <sub>2</sub> SO <sub>4</sub> ; CH <sub>3</sub> Cl/AlCl <sub>3</sub> ; KMnO <sub>4</sub> /OH <sup>-</sup> (C) CH <sub>3</sub> Cl/AlCl <sub>3</sub> ; KMnO <sub>4</sub> /OH <sup>-</sup> ; HNO <sub>3</sub> /H <sub>2</sub> SO <sub>4</sub> (D) CH <sub>3</sub> Cl/AlCl <sub>3</sub> ; HNO <sub>3</sub> /H <sub>2</sub> SO <sub>4</sub> ; KMnO <sub>4</sub> /OH <sup>-</sup>				
Ans.	(D)				
Sol.	$(1) CH_3CI$ $(2) HNO_3/H_2SO_4$ $(3) KMnO_4/OH^-$ $(3) KMnO_4/OH^-$ $(1) CH_3CI$ $(2) HNO_3/H_2SO_4$ $(3) KMnO_4/OH^-$				
57. Ans	The volume of one drop of aqueous solution from an eyedropper is approximately 0.05 mL. One such drop of 0.2 M HCl is added to 100 mL of distilled water. The pH of the resulting solution will be (A) 4.0 (B) 7.0 (C) 3.0 (D) 5.5				
AII5.	(2) 0.2×0.05				
Sol.	Conc. of HCl = $\frac{10^{-4} \text{ M}}{100}$ = 10 <sup>-4</sup> M				
	50 pπ = 4				
58. Ans.	In which of the following species the octet rule is NOT obeyed? I. $I_3^-$ II. $N_2O$ III. $OF_2$ IV. $NO^+$ (A) I and IV (B) II and III (C) I only (D) IV only (C)				
Sol.	$I_3^ [I_3^-, I]^-$ Octet rule not obeyed				
	N2Ofollowsoctet ruleOF2followsoctet ruleNO+followsoctet rule				
59.	Which atom/s will have a $\delta^+$ charge in the following molecule ?				
Ans.	(A) I and III (B) II only (C) II and III (D) II and IV (D)				
60.	2.0 moles of an ideal gas expands isothermally (27°C) and reversibly from a pressure of 1 bar to 10 bar. The heaviest mass that can be lifted through a height of 10 m by the work of this expansion is (A) 50.8 kg (B) 50.8 g (C) 117.1 kg (D) 117.1 g				
Ans.	(C) Resonance® Reg. & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.)- 324005 Website : www.resonance.ac.in   E-mail : contact@resonance.ac.in				
	Constraint         Constra				

Sol.	work done  = change in potential energy							
	$nRTIn\frac{V_2}{V_1} = mgh$							
	$2 \times 8.314 \times 300 \times 2.303 \log \frac{10}{1} = m \times 9.81 \times 10$							
	m = 1	117.1 kg	·					
	In Q. Nos. 6 identify all o correct answ	1 to 70 any f them cor wer is miss	/ number of option rectly to get 6 ma red, you get zero	A – 2 ons (A arks. E marks	or B or C or ven if one ans	all D) ma swer ider	ay be corre ntified is in	ect. You are to correct or one
61.	A commercia percentage o (A) 2.88	Il sample of If free SO₃ i	oleum (H <sub>2</sub> S <sub>2</sub> O <sub>7</sub> ) la n this oleum samp (B) 28.8	abeled ole is	as '106.5% ole (C) 0.029	um' cont	ains 6.5 g c (D) 0.28	of water. The
Ans.	(B)							
501.	106.5%	Oleum s	sample					
	∴ % of free S	$SO_3 = \frac{40}{9} \times$	(106.5–100) = 28	8.8 %				
	Note: Questi mixture. Also	on is contro oleum doe	overcial since forn s not contains wa	nula of ter, hov	oleum is giver vever in questio	n (H₂S₂O on, it is sa	7), however aid to be pr	it is actually a esent.
62.	Which of the	following sp	becies has one lor	ne pair	of electrons on	the cent	ral atom?	
Ans.	(A) CIF₃ (D) F		(B) I <sub>3</sub> -		(C) I₃⁺ F		(D) SF4	
Sol.	[CI—F   F	[I <b>Ï</b>	ul- I ↓	I	F			
63.	Among the fo I. [Ni(CO)4] (A) I and IV	bllowing, the	e complex ion/s th II. [NiCl₄] <sup>2–</sup> (B) II only	at will h	nave a magneti III. [Ni(H₂O)₀] <sup>2+</sup> (C) II and III	c momen	t of 2.82 B. IV. [Ni(CN (D) II, III a	M. is/are )₄] <sup>2–</sup> nd IV
Ans. Sol.	(c) Ni(CO) <sub>4</sub> number of unpaired electron = 0 [NiCl <sub>4</sub> ] <sup>2-</sup> number of unpaired electron = 2 [Ni(H <sub>2</sub> O) <sub>6</sub> ] <sup>2+</sup> number of unpaired electron = 2 [Ni(CN) <sub>4</sub> ] <sup>2-</sup> number of unpaired electron = 0 $\mu = \sqrt{n(n+2)} \Rightarrow 2.82 = \sqrt{n(n+2)}$ n = number of unpaired electrons = 2							
64.	Morphine, a pain killer is basic with molecular formula $C_{17}H_{19}NO_3$ . The conjugate acid of morphine							
Ans.	(A) C <sub>17</sub> H <sub>19</sub> NC (D)	<b>)</b> <sub>3</sub> +	(B) C <sub>17</sub> H <sub>18</sub> NO <sub>3</sub>		(C) C <sub>17</sub> H <sub>19</sub> NO <sub>3</sub>	-	(D) C <sub>17</sub> H <sub>20</sub>	NO <sub>3</sub> +
65.	A suboxide of carbon, $C_3O_2$ , has a linear structure. Which of the following is correct about $C_3O_2$ ? I. Oxidation state of all three C atoms is +2 II. Oxidation state of the central C atom is zero III. The molecule contains 4 $\sigma$ and 4 $\pi$ bonds IV. Hybridization of the central carbon atom is sp <sup>2</sup> (A) based by (C) if a bit (C) if							
Ans.	(A) I and IV (B)		(B) II and III	-	(C) II and IV		(D) III only	
Sol.	Hyb. Sp s	0 +2 C=C=O sp sp						
		®	Reg. & Corp. Office: C	G Tower,	A-46 & 52, IPIA, Nea	ar City Mall,	Jhalawar Road,	Kota (Raj.)- 324005
$\square$	Keson Educating for bet		Website : www.resonan	nce.ac.in	E-mail : contact@res	sonance.ac.ir	NS	EC24112019 # 20
	Toll Free : 1800 258 5555   CIN: U80302RJ2007PLC024029         COLORATION # 20							







The reaction is

- (A) exothermic and K<sub>eq</sub> at 1400 K =  $3.79 \times 10^{-6}$
- (B) endothermic and K<sub>eq</sub> at 1400 K =  $2.63 \times 10^{-5}$
- (C) exothermic and  $K_{eq}$  at 1400 K = 1.8 × 10<sup>5</sup>
- (D) endothermic and K<sub>eq</sub> at 1500 K =  $9.28 \times 10^{-4}$

**Sol.** At 1400 K

$$K_{eq} = \frac{K_{f}}{R_{b}} = \frac{0.2}{1.1 \times 10^{-6}} = 1.8 \times 10^{5}$$

At 1500 K

 $K_{eq} = \frac{1.3}{1.4 \times 10^{-5}} = 9.3 \times 10^{4}$ 

: reaction is exothermic



ß	Reg. & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar R	oad, Kota (Raj.)- 324005
sonance	Website : www.resonance.ac.in   E-mail : contact@resonance.ac.in	NSEC24112010 # 22
ng for better tomorrow	Toll Free : 1800 258 5555   CIN: U80302RJ2007PLC024029	N3EC24112019 # 23



INATIONAL STANDARD EXAMINATION IN CHEMISTRY (NSEC) 2019-20 | 24-11-2019



Resonance®





# **ADMISSION OPEN**

For Classes: V to XII & XII+ (XII Passed)

# JEE ADVANCED | JEE MAIN PRE-MEDICAL | COMMERCE & LAW PCCP (V to X) | OLYMPIADS NTSE | IJSO | KVPY

Enroll Now for Academic Session 2020-21 @ Coaching Fee of 2019-20

# Call Now: 1800 258 5555

# Academic Benefits\*

# **Financial Benefits\***

- More than 800 Academic Hours
- More than 500 Classes
- More than 100 Testing Hours

**ResoFAST Test Dates:** 08<sup>th</sup>, 22<sup>nd</sup>, & 29<sup>th</sup> Dec 2019

# • Upto Rs.35000+ Saving on 1 Year Course Fee

50% Concession on Admission Form Fee

L& CApply

• Up to 90% Scholarship on Course Fee

**Resonance Eduventures Limited** Registered & Corporate Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Rajasthan) - 324005 | Tel.: 0744-2777777, 2777700 | CIN: U80302RJ2007PLC024029

Website: www.resonance.ac.in | e-mail: contact@resonance.ac.in | f 💌 🛅 🕒



 Reg. & Corp. Office: CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.)- 324005

 Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in

 Toll Free : 1800 258 5555 | CIN: U80302RJ2007PLC024029

NSEC24112019 # 25