

## PART I : PHYSICS

## SECTION I : Single Correct Answer Type

This section contains 10 multiple choice questions. Each question has four choices (A), (B), (C) and (D) out of which **ONLY ONE** is correct.


1. Two large vertical and parallel metal plates having a separation of 1 cm are connected to a DC voltage source of potential difference  $X$ . A proton is released at rest midway between the two plates. It is found to move at  $45^\circ$  to the vertical JUST after release. Then  $X$  is nearly  
 (A)  $1 \times 10^{-5}$  V       (B)  $1 \times 10^{-7}$  V      (C)  $1 \times 10^{-9}$  V      (D)  $1 \times 10^{-10}$  V

2. A mixture of 2 moles of helium gas (atomic mass = 4 amu) and 1 mole of argon gas (atomic mass = 40 amu) is kept at 300 K in a container. The ratio of the rms speeds

$$\left( \frac{v_{rms}(\text{helium})}{v_{rms}(\text{argon})} \right) \text{ is}$$

- (A) 0.32      (B) 0.45      (C) 2.24       (D) 3.16

3. A small block is connected to one end of a massless spring of un-stretched length 4.9 m. The other end of the spring (see the figure) is fixed. The system lies on a horizontal frictionless surface. The block is stretched by 0.2 m and released from rest at  $t = 0$ . It then executes simple harmonic motion with angular frequency  $\omega = \frac{\pi}{3}$  rad/s. Simultaneously at  $t = 0$ , a small pebble is projected with speed  $v$  from point  $P$  at an angle of  $45^\circ$  as shown in the figure. Point  $P$  is at a horizontal distance of 10 m from  $O$ . If the pebble hits the block at  $t = 1$  s, the value of  $v$  is (take  $g = 10 \text{ m/s}^2$ )




- (A)  $\sqrt{50}$  m/s      (B)  $\sqrt{51}$  m/s      (C)  $\sqrt{52}$  m/s      (D)  $\sqrt{53}$  m/s

Space for Rough Work

4. Three very large plates of same area are kept parallel and close to each other. They are considered as ideal black surfaces and have very high thermal conductivity. The first and third plates are maintained at temperatures  $2T$  and  $3T$  respectively. The temperature of the middle (i.e. second) plate under steady state condition is

- (A)  $\left(\frac{65}{2}\right)^{\frac{1}{4}} T$       (B)  $\left(\frac{97}{4}\right)^{\frac{1}{4}} T$        (C)  $\left(\frac{97}{2}\right)^{\frac{1}{4}} T$       (D)  $(97)^{\frac{1}{4}} T$

5. A thin uniform rod, pivoted at  $O$ , is rotating in the horizontal plane with constant angular speed  $\omega$ , as shown in the figure. At time  $t = 0$ , a small insect starts from  $O$  and moves with constant speed  $v$  with respect to the rod towards the other end. It reaches the end of the rod at  $t = T$  and stops. The angular speed of the system remains  $\omega$  throughout. The magnitude of the torque ( $|\vec{\tau}|$ ) on the system about  $O$ , as a function of time is best represented by which plot ?



- (A)      (B)      (C)       (D)

Space for Rough Work

Handwritten notes and diagrams for problem 4 and 5. For problem 4, temperatures  $2T$  and  $3T$  are noted, and calculations involving  $\frac{27+3T}{2}$  and  $\left(\frac{97}{2}\right)^{\frac{1}{4}}$  are shown. For problem 5, a diagram of the rod and insect is repeated, and calculations for the torque  $\tau = r \times mv$  are shown, leading to the conclusion that the torque increases with a concave-up curve.


6. In the determination of Young's modulus  $\left( Y = \frac{4MLg}{\pi d^2 l} \right)$  by using Searle's method, a wire of length  $L = 2$  m and diameter  $d = 0.5$  mm is used. For a load  $M = 2.5$  kg, an extension  $l = 0.25$  mm in the length of the wire is observed. Quantities  $d$  and  $l$  are measured using a screw gauge and a micrometer, respectively. They have the same pitch of 0.5 mm. The number of divisions on their circular scale is 100. The contributions to the maximum probable error of the  $Y$  measurement

- (A) due to the errors in the measurements of  $d$  and  $l$  are the same.
- ✓ (B) due to the error in the measurement of  $d$  is twice that due to the error in the measurement of  $l$ .
- (C) due to the error in the measurement of  $l$  is twice that due to the error in the measurement of  $d$ .
- (D) due to the error in the measurement of  $d$  is four times that due to the error in the measurement of  $l$ .

Space for Rough Work

$Y = \frac{4MLg}{\pi d^2 l}$   
 $L = 2$  m     $d = 0.5$  mm  
 $M = 2.5$  kg  
 $l = 0.25$  mm  
 pitch = 0.5 mm  
 no. of div. = 100  
 error =  $\frac{0.5}{100} = 0.005$  mm  
 ✓

7. Consider a thin spherical shell of radius  $R$  with its centre at the origin, carrying uniform positive surface charge density. The variation of the magnitude of the electric field  $|\vec{E}(r)|$  and the electric potential  $V(r)$  with the distance  $r$  from the centre, is best represented by which graph ?




Space for Rough Work

Handwritten notes and diagrams for rough work:

- Labels:  $R$ ,  $|\vec{E}(r)|$ ,  $V(r)$ ,  $r$ .
- Equation:  $V = \int \vec{E} \cdot d\vec{r}$
- Equation:  $E = \frac{V}{r}$
- Diagram: A sketch of a spherical shell with radius  $R$  and a point at distance  $r$  from the center. Electric field lines are shown as arrows pointing outwards from the shell.



8. A bi-convex lens is formed with two thin plano-convex lenses as shown in the figure. Refractive index  $n$  of the first lens is 1.5 and that of the second lens is 1.2. Both the curved surfaces are of the same radius of curvature  $R = 14$  cm. For this bi-convex lens, for an object distance of 40 cm, the image distance will be



- (A) - 280.0 cm.      (B) 40.0 cm.      (C) 21.5 cm.      (D) 13.3 cm.

Space for Rough Work


$n_1 = 1.5$   
 $n_2 = 1.2$   
 $R = 14 \text{ cm}$

$f_1 = \frac{R}{n_1 - 1} = \frac{14}{1.5 - 1} = \frac{14}{0.5} = 28 \text{ cm}$   
 $f_2 = \frac{R}{n_2 - 1} = \frac{14}{1.2 - 1} = \frac{14}{0.2} = 70 \text{ cm}$

$\frac{1}{f} = \frac{1}{f_1} + \frac{1}{f_2}$   
 $\frac{1}{f} = \frac{1}{28} + \frac{1}{70}$   
 $\frac{1}{f} = \frac{5}{140} + \frac{2}{140} = \frac{7}{140} = \frac{1}{20}$   
 $f = 20 \text{ cm}$

For an object distance  $u = 40 \text{ cm}$ ,  
 $\frac{1}{v} - \frac{1}{u} = \frac{1}{f}$   
 $\frac{1}{v} - \frac{1}{40} = \frac{1}{20}$   
 $\frac{1}{v} = \frac{1}{20} + \frac{1}{40} = \frac{2}{40} + \frac{1}{40} = \frac{3}{40}$   
 $v = \frac{40}{3} \approx 13.3 \text{ cm}$

9. A small mass  $m$  is attached to a massless string whose other end is fixed at  $P$  as shown in the figure. The mass is undergoing circular motion in the  $x$ - $y$  plane with centre at  $O$  and constant angular speed  $\omega$ . If the angular momentum of the system, calculated about  $O$  and  $P$  are denoted by  $\vec{L}_O$  and  $\vec{L}_P$  respectively, then



- (A)  $\vec{L}_O$  and  $\vec{L}_P$  do not vary with time.
- (B)  $\vec{L}_O$  varies with time while  $\vec{L}_P$  remains constant.
- (C)  $\vec{L}_O$  remains constant while  $\vec{L}_P$  varies with time.
- (D)  $\vec{L}_O$  and  $\vec{L}_P$  both vary with time.

10. Young's double slit experiment is carried out by using green, red and blue light, one color at a time. The fringe widths recorded are  $\beta_G, \beta_R$  and  $\beta_B$ , respectively. Then,

- (A)  $\beta_G > \beta_B > \beta_R$
- (B)  $\beta_B > \beta_G > \beta_R$
- (C)  $\beta_R > \beta_B > \beta_G$
- (D)  $\beta_R > \beta_G > \beta_B$

Space for Rough Work

*Handwritten notes and diagrams:*

$m$   $P$

$x$ - $y$  plane  $O$   
angular speed  $\omega$

$L_O$   $L_P$

$\lambda$   $D_s$   $G$   $R$   $B$

1 color at a time


fringe width  $\beta_G > \beta_R > \beta_B$



## SECTION II : Multiple Correct Answer(s) Type

This section contains 5 multiple choice questions. Each question has four choices (A), (B), (C) and (D) out of which ONE or MORE are correct.

11. A person blows into open-end of a long pipe. As a result, a high-pressure pulse of air travels down the pipe. When this pulse reaches the other end of the pipe,
- (A) a high-pressure pulse starts traveling up the pipe, if the other end of the pipe is open.  
 (B) a low-pressure pulse starts traveling up the pipe, if the other end of the pipe is open.  
 (C) a low-pressure pulse starts traveling up the pipe, if the other end of the pipe is closed.  
 (D) a high-pressure pulse starts traveling up the pipe, if the other end of the pipe is closed.
12. A small block of mass of 0.1 kg lies on a fixed inclined plane  $PQ$  which makes an angle  $\theta$  with the horizontal. A horizontal force of 1 N acts on the block through its center of mass as shown in the figure. The block remains stationary if (take  $g = 10 \text{ m/s}^2$ )



- (A)  $\theta = 45^\circ$ .  
 (B)  $\theta > 45^\circ$  and a frictional force acts on the block towards  $P$ .  
 (C)  $\theta > 45^\circ$  and a frictional force acts on the block towards  $Q$ .  
 (D)  $\theta < 45^\circ$  and a frictional force acts on the block towards  $Q$ .

Space for Rough Work

Handwritten rough work for Question 12:

Diagram of the inclined plane with a block. A horizontal force of 1 N is applied to the block. The mass of the block is 0.1 kg. The angle of the incline is  $\theta$ . The acceleration due to gravity is  $g = 10 \text{ m/s}^2$ .


Handwritten notes:

$\mu = 0.5$   
 $P_f = mg = 10 \times 0.1 \times 10$

Page No. : 9



13. A cubical region of side  $a$  has its centre at the origin. It encloses three fixed point charges,  $-q$  at  $(0, -a/4, 0)$ ,  $+3q$  at  $(0, 0, 0)$  and  $-q$  at  $(0, +a/4, 0)$ . Choose the correct option(s).



- (A) The net electric flux crossing the plane  $x = +a/2$  is equal to the net electric flux crossing the plane  $x = -a/2$ .
- (B) The net electric flux crossing the plane  $y = +a/2$  is more than the net electric flux crossing the plane  $y = -a/2$ .
- (C) The net electric flux crossing the entire region is  $\frac{q}{\epsilon_0}$ .
- (D) The net electric flux crossing the plane  $z = +a/2$  is equal to the net electric flux crossing the plane  $x = +a/2$ .

Space for Rough Work

$-q$  at  $(0, -a/4, 0)$   
 $+3q$  at  $(0, 0, 0)$   
 $-q$  at  $(0, +a/4, 0)$

flux = 0







SECTION III : Integer Answer Type



This section contains 5 questions. The answer to each question is a single digit integer, ranging from 0 to 9 (both inclusive).

16. A proton is fired from very far away towards a nucleus with charge  $Q = 120 e$ , where  $e$  is the electronic charge. It makes a closest approach of 10 fm to the nucleus. The de Broglie wavelength (in units of fm) of the proton at its start is: (take the proton mass,  $m_p = (5/3) \times 10^{-27}$  kg;  $h/e = 4.2 \times 10^{-15}$  J.s/C;  $\frac{1}{4\pi\epsilon_0} = 9 \times 10^9$  m/F;  $1 \text{ fm} = 10^{-15}$  m) 6

17. A lamina is made by removing a small disc of diameter  $2R$  from a bigger disc of uniform mass density and radius  $2R$ , as shown in the figure. The moment of inertia of this lamina about axes passing through  $O$  and  $P$  is  $I_O$  and  $I_P$ , respectively. Both these axes are perpendicular to the plane of the lamina. The ratio  $\frac{I_P}{I_O}$  to the nearest integer is



18. An infinitely long solid cylinder of radius  $R$  has a uniform volume charge density  $\rho$ . It has a spherical cavity of radius  $R/2$  with its centre on the axis of the cylinder, as shown in the figure. The magnitude of the electric field at the point  $P$ , which is at a distance  $2R$  from the axis of the cylinder, is given by the expression  $\frac{23\rho R}{16k\epsilon_0}$ . The value of  $k$  is



Space for Rough Work

\*4



$Q = 120e$   
 $m_p = 5/3 \times 10^{-27}$

$\frac{1}{4\pi\epsilon_0} = 9 \times 10^9$



$h/e = 4.2 \times 10^{-15}$

19. A cylindrical cavity of diameter  $a$  exists inside a cylinder of diameter  $2a$  as shown in the figure. Both the cylinder and the cavity are infinitely long. A uniform current density  $J$  flows along the length.

If the magnitude of the magnetic field at the point  $P$  is given by  $\frac{N}{12} \mu_0 a J$ , then the value of  $N$  is




20. A circular wire loop of radius  $R$  is placed in the  $x$ - $y$  plane centered at the origin  $O$ . A square loop of side  $a$  ( $a \ll R$ ) having two turns is placed with its center at  $z = \sqrt{3} R$  along the axis of the circular wire loop, as shown in figure. The plane of the square loop makes an angle of  $45^\circ$  with respect to the  $z$ -axis. If the mutual inductance between the loops is given by  $\frac{\mu_0 a^2}{2^{p/2} R}$ , then the value of  $p$  is



Space for Rough Work



24. For one mole of a van der Waals gas when  $b = 0$  and  $T = 300$  K, the  $PV$  vs.  $1/V$  plot is shown below. The value of the van der Waals constant  $a$  ( $\text{atm.liter}^2 \text{mol}^{-2}$ ) is



$b = 0$   $T = 300$   
 $PV$  vs  $1/V$   
 Van der Waals (1 mole (1/mole<sup>2</sup>))

(A) 1.0

(B) 4.5

(C) 1.5

(D) 3.0

25. In allene ( $\text{C}_3\text{H}_4$ ), the type(s) of hybridisation of the carbon atoms is (are)


(A)  $sp$  and  $sp^3$

(B)  $sp$  and  $sp^2$

(C) only  $sp^2$

(D)  $sp^2$  and  $sp^3$

26. A compound  $\text{M}_p\text{X}_q$  has cubic close packing (ccp) arrangement of X. Its unit cell structure is shown below. The empirical formula of the compound is



M =  $\square$   
 X =  $\circ$

(A)  $\text{MX}$

(B)  $\text{MX}_2$

(C)  $\text{M}_2\text{X}$


(D)  $\text{M}_5\text{X}_{14}$

$\text{C}_3\text{H}_4$   
 $\text{C} + 2\text{O}_2 \rightarrow \text{CO}_2$   
 $\text{M}_2 \text{ M}_4$   
 CCP  
 $2 + 25$   
 $\text{C}_3\text{H}_4\text{O}_2$

CHEMISTRY

Space for Rough Work

27. The number of optically active products obtained from the complete ozonolysis of the given compound is,



- (A) 0                      (B) 1                       (C) 2                      (D) 4

28. As per IUPAC nomenclature, the name of the complex  $[\text{Co}(\text{H}_2\text{O})_4(\text{NH}_3)_2]\text{Cl}_3$  is

- (A) Tetraaquadiaminecobalt (III) chloride  
 (B) Tetraaquadiamminecobalt (III) chloride  
 (C) Diaminetetraaquacobalt (III) chloride  
 (D) Diamminetetraaquacobalt (III) chloride



29. The carboxyl functional group ( $-\text{COOH}$ ) is present in

- (A) picric acid              (B) barbituric acid              (C) ascorbic acid              (D) aspirin

30. The colour of light absorbed by an aqueous solution of  $\text{CuSO}_4$  is

- (A) orange-red               (B) blue-green              (C) yellow              (D) violet

Space for Rough Work




(A)

SECTION II : Multiple Correct Answer(s) Type

This section contains 5 multiple choice questions. Each question has four choices (A), (B), (C) and (D) out of which ONE or MORE are correct.

31. For an ideal gas, consider only  $P$ - $V$  work in going from an initial state  $X$  to the final state  $Z$ . The final state  $Z$  can be reached by either of the two paths shown in the figure. Which of the following choice(s) is (are) correct? [take  $\Delta S$  as change in entropy and  $w$  as work done]







(A)  $\Delta S_{x \rightarrow z} = \Delta S_{x \rightarrow y} + \Delta S_{y \rightarrow z}$

(B)  $w_{x \rightarrow z} = w_{x \rightarrow y} + w_{y \rightarrow z}$



(C)  $w_{x \rightarrow y \rightarrow z} = w_{x \rightarrow y}$

(D)  $\Delta S_{x \rightarrow y \rightarrow z} = \Delta S_{x \rightarrow y}$


32. Which of the following molecules, in pure form, is (are) **unstable** at room temperature?



Space for Rough Work



33. Identify the binary mixture(s) that can be separated into individual compounds, by differential extraction, as shown in the given scheme.



- (A)  $C_6H_5OH$  and  $C_6H_5COOH$        (B)  $C_6H_5COOH$  and  $C_6H_5CH_2OH$   
 (C)  $C_6H_5CH_2OH$  and  $C_6H_5OH$       (D)  $C_6H_5CH_2OH$  and  $C_6H_5CH_2COOH$


34. Choose the correct reason(s) for the stability of the **lyophobic** colloidal particles.

- (A) Preferential adsorption of ions on their surface from the solution  
 (B) Preferential adsorption of solvent on their surface from the solution  
 (C) Attraction between different particles having opposite charges on their surface  
 (D) Potential difference between the fixed layer and the diffused layer of opposite charges around the colloidal particles

35. Which of the following hydrogen halides react(s) with  $AgNO_3(aq)$  to give a precipitate that dissolves in  $Na_2S_2O_3(aq)$  ?

- (A) HCl       (B) HF       (C) HBr      (D) HI

Space for Rough Work





SECTION III : Integer Answer Type


This section contains 5 questions. The answer to each question is a single digit integer, ranging from 0 to 9 (both inclusive).

36. An organic compound undergoes first-order decomposition. The time taken for its decomposition to 1/8 and 1/10 of its initial concentration are  $t_{1/8}$  and  $t_{1/10}$  respectively.

What is the value of  $\frac{[t_{1/8}]}{[t_{1/10}]} \times 10$ ? (take  $\log_{10} 2 = 0.3$ )

3

37. When the following aldohexose exists in its D-configuration, the total number of stereoisomers in its pyranose form is



5

CHEMISTRY

Space for Rough Work

1<sup>st</sup> order dec<sup>n</sup>


$T = \frac{1}{k} \ln \frac{1}{2}$

$t_{1/8} \quad t_{1/10}$

$\frac{[t_{1/8}]}{[t_{1/10}]} = \frac{\ln 8}{\ln 10}$


$\log_{10} 2 = 0.3$

38. The substituents  $R_1$  and  $R_2$  for nine peptides are listed in the table given below. How many of these peptides are positively charged at  $\text{pH} = 7.0$  ?



Peptide	$R_1$	$R_2$
I	H	H
II	H	$\text{CH}_3$
III	$\text{CH}_2\text{COOH}$	H
IV	$\text{CH}_2\text{CONH}_2$	$(\text{CH}_2)_4\text{NH}_2$
V	$\text{CH}_2\text{CONH}_2$	$\text{CH}_2\text{CONH}_2$
VI	$(\text{CH}_2)_4\text{NH}_2$	$(\text{CH}_2)_4\text{NH}_2$
VII	$\text{CH}_2\text{COOH}$	$\text{CH}_2\text{CONH}_2$
VIII	$\text{CH}_2\text{OH}$	$(\text{CH}_2)_4\text{NH}_2$
IX	$(\text{CH}_2)_4\text{NH}_2$	$\text{CH}_3$

39. The periodic table consists of 18 groups. An isotope of copper, on bombardment with protons, undergoes a nuclear reaction yielding element  $X$  as shown below. To which group, element  $X$  belongs in the periodic table ?



40. 29.2% (w/w) HCl stock solution has a density of  $1.25 \text{ g mL}^{-1}$ . The molecular weight of HCl is  $36.5 \text{ g mol}^{-1}$ . The volume (mL) of stock solution required to prepare a 200 mL solution of 0.4 M HCl is

Space for Rough Work

$29.2\%$  (w/w) HCl soln  $d = 1.25 \text{ g mL}^{-1}$   
 $M.W. \text{ HCl} = 36.5 \text{ g mol}^{-1}$   
 $200 \text{ mL soln } 0.4 \text{ M HCl}$

# PART III : MATHEMATICS

## SECTION I : Single Correct Answer Type

This section contains 10 multiple choice questions. Each question has four choices (A), (B), (C) and (D) out of which **ONLY ONE** is correct.

41. The point  $P$  is the intersection of the straight line joining the points  $Q(2,3,5)$  and  $R(1, -1, 4)$  with the plane  $5x - 4y - z = 1$ . If  $S$  is the foot of the perpendicular drawn from the point  $T(2, 1, 4)$  to  $QR$ , then the length of the line segment  $PS$  is

- (A)  $\frac{1}{\sqrt{2}}$                       (B)  $\sqrt{2}$                       (C) 2                      (D)  $2\sqrt{2}$

42. The integral  $\int \frac{\sec^2 x}{(\sec x + \tan x)^{3/2}} dx$  equals (for some arbitrary constant  $K$ )

(A)  $-\frac{1}{(\sec x + \tan x)^{1/2}} \left\{ \frac{1}{11} - \frac{1}{7} (\sec x + \tan x)^2 \right\} + K$

(B)  $\frac{1}{(\sec x + \tan x)^{1/2}} \left\{ \frac{1}{11} - \frac{1}{7} (\sec x + \tan x)^2 \right\} + K$

(C)  $-\frac{1}{(\sec x + \tan x)^{1/2}} \left\{ \frac{1}{11} + \frac{1}{7} (\sec x + \tan x)^2 \right\} + K$

(D)  $\frac{1}{(\sec x + \tan x)^{1/2}} \left\{ \frac{1}{11} + \frac{1}{7} (\sec x + \tan x)^2 \right\} + K$

Space for Rough Work

*Handwritten notes for Q41:*

$Q(2,3,5)$   
 $R(1,-1,4)$   
 $5x - 4y - z = 1$   
 $T(2,1,4)$   
 $PS$

*Handwritten notes for Q42:*

$\int \frac{\sec^2 x}{(\sec x + \tan x)^{3/2}} dx$   
 $\frac{1}{\sec x + \tan x}$



47. The function  $f: [0, 3] \rightarrow [1, 29]$ , defined by  $f(x) = 2x^3 - 15x^2 + 36x + 1$ , is

- (A) one-one and onto.
- (B) onto but not one-one.
- (C) one-one but not onto.
- (D) neither one-one nor onto.

48. The locus of the mid-point of the chord of contact of tangents drawn from points lying on the straight line  $4x - 5y = 20$  to the circle  $x^2 + y^2 = 9$  is

- (A)  $20(x^2 + y^2) - 36x + 45y = 0$
- (B)  $20(x^2 + y^2) + 36x - 45y = 0$
- (C)  $36(x^2 + y^2) - 20x + 45y = 0$
- (D)  $36(x^2 + y^2) + 20x - 45y = 0$

49. Let  $P = [a_{ij}]$  be a  $3 \times 3$  matrix and let  $Q = [b_{ij}]$ , where  $b_{ij} = 2^{i+j}a_{ij}$  for  $1 \leq i, j \leq 3$ . If the determinant of  $P$  is 2, then the determinant of the matrix  $Q$  is

- (A)  $2^{10}$
- (B)  $2^{11}$
- (C)  $2^{12}$
- (D)  $2^{13}$

50. The ellipse  $E_1: \frac{x^2}{9} + \frac{y^2}{4} = 1$  is inscribed in a rectangle  $R$  whose sides are parallel to the coordinate axes. Another ellipse  $E_2$  passing through the point  $(0, 4)$  circumscribes the rectangle  $R$ . The eccentricity of the ellipse  $E_2$  is

- (A)  $\frac{\sqrt{2}}{2}$
- (B)  $\frac{\sqrt{3}}{2}$
- (C)  $\frac{1}{2}$
- (D)  $\frac{3}{4}$

$P = [a_{ij}]$   $3 \times 3$   
 $Q = [b_{ij}]$   $3 \times 3$   
 $b_{ij} = 2^{i+j} a_{ij}$

Space for Rough Work

$4x - 5y = 20$   $1 \leq i, j \leq 3$   
 $x^2 + y^2 = 9$   $P = 2$   
 $x \in \mathbb{R}$   $3 \times 3$   
 $20x^2 + 45y^2 = 0$   
 $\frac{x^2 + x - 1}{x - 1} = 2$

$f: [0, 3] \rightarrow [1, 29]$   $f(x) = 2x^3 - 15x^2 + 36x + 1$   
 one-one ✓



53. Let  $\theta, \varphi \in [0, 2\pi]$  be such that

$$2 \cos \theta (1 - \sin \varphi) = \sin^2 \theta \left( \tan \frac{\theta}{2} + \cot \frac{\theta}{2} \right) \cos \varphi - 1,$$

$$\tan(2\pi - \theta) > 0 \text{ and } -1 < \sin \theta < -\frac{\sqrt{3}}{2}.$$

Then  $\varphi$  cannot satisfy

(A)  $0 < \varphi < \frac{\pi}{2}$

(B)  $\frac{\pi}{2} < \varphi < \frac{4\pi}{3}$

(C)  $\frac{4\pi}{3} < \varphi < \frac{3\pi}{2}$

(D)  $\frac{3\pi}{2} < \varphi < 2\pi$

54. Let  $S$  be the area of the region enclosed by  $y = e^{-x^2}$ ,  $y = 0$ ,  $x = 0$ , and  $x = 1$ . Then

(A)  $S \geq \frac{1}{e}$

(B)  $S \geq 1 - \frac{1}{e}$

(C)  $S \leq \frac{1}{4} \left( 1 + \frac{1}{\sqrt{e}} \right)$

(D)  $S \leq \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{e}} \left( 1 - \frac{1}{\sqrt{2}} \right)$

55. Tangents are drawn to the hyperbola  $\frac{x^2}{9} - \frac{y^2}{4} = 1$ , parallel to the straight line  $2x - y = 1$ . The points of contact of the tangents on the hyperbola are

(A)  $\left( \frac{9}{2\sqrt{2}}, \frac{1}{\sqrt{2}} \right)$

(B)  $\left( -\frac{9}{2\sqrt{2}}, -\frac{1}{\sqrt{2}} \right)$

(C)  $(3\sqrt{3}, -2\sqrt{2})$

(D)  $(-3\sqrt{3}, 2\sqrt{2})$

Space for Rough Work

$\varphi \in [0, 2\pi]$

$2 \cos \theta (1 - \sin \varphi) = \sin^2 \theta \left( \tan \frac{\theta}{2} + \cot \frac{\theta}{2} \right) \cos \varphi - 1$

$\tan(2\pi - \theta) > 0$  &  $-1 < \sin \theta < -\frac{\sqrt{3}}{2}$

$S$   $y = e^{-x^2}$ ,  $y = 0$ ,  $x = 0$  &  $x = 1$

SECTION III : Integer Answer Type

This section contains 5 questions. The answer to each question is a single digit integer, ranging from 0 to 9 (both inclusive).

56. Let  $S$  be the focus of the parabola  $y^2 = 8x$  and let  $PQ$  be the common chord of the circle  $x^2 + y^2 - 2x - 4y = 0$  and the given parabola. The area of the triangle  $PQS$  is 2

57. Let  $p(x)$  be a real polynomial of least degree which has a local maximum at  $x = 1$  and a local minimum at  $x = 3$ . If  $p(1) = 6$  and  $p(3) = 2$ , then  $p'(0)$  is 0

58. Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be defined as  $f(x) = |x| + |x^2 - 1|$ . The total number of points at which  $f$  attains either a local maximum or a local minimum is 1

59. The value of  $6 + \log_{\frac{3}{2}} \left( \frac{1}{3\sqrt{2}} \sqrt{4 - \frac{1}{3\sqrt{2}} \sqrt{4 - \frac{1}{3\sqrt{2}} \sqrt{4 - \frac{1}{3\sqrt{2}} \dots}}} \right)$  is 4

60. If  $\vec{a}, \vec{b}$  and  $\vec{c}$  are unit vectors satisfying  $|\vec{a} - \vec{b}|^2 + |\vec{b} - \vec{c}|^2 + |\vec{c} - \vec{a}|^2 = 9$ , then  $|2\vec{a} + 5\vec{b} + 5\vec{c}|$  is 6

Space for Rough Work

$\frac{x^2}{9} - \frac{y^2}{4} = 1$        $y^2 = 8x$   
 $2x - y = 1$        $PQ$  common chord  
 $x^2 + y^2 - 2x - 4y = 0$   
 Area  $\Delta PQS$   
 w/wc.  $x = 1$       w/wc.  $x = 3$   
 $p(1) = 6$        $p(3) = 2$   
 $p'(0)$   
 $f: \mathbb{R} \rightarrow \mathbb{R} \rightarrow f(x) = |x| + |x^2 - 1|$