



**Resonance**<sup>®</sup>  
Educating for better tomorrow

# JEE

## (Main)

**PAPER-1 (B.E./B. TECH.)**

# 2022

## **COMPUTER BASED TEST (CBT)** **Questions & Solutions**

**Date: 29 July, 2022 (SHIFT-2) | TIME : (3.00 a.m. to 6.00 p.m)**

**Duration: 3 Hours | Max. Marks: 300**






**SUBJECT: PHYSICS**

### **Resonance Eduventures Ltd.**

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : [www.resonance.ac.in](http://www.resonance.ac.in) | E-mail : [contact@resonance.ac.in](mailto:contact@resonance.ac.in) | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555  7340010333  [facebook.com/ResonanceEdu](https://facebook.com/ResonanceEdu)  [twitter.com/ResonanceEdu](https://twitter.com/ResonanceEdu)  [www.youtube.com/resowatch](https://www.youtube.com/resowatch)  [blog.resonance.ac.in](https://blog.resonance.ac.in)

This solution was download from Resonance JEE (MAIN) 2022 Solution portal

**PART : PHYSICS**

1. Two identical metallic spheres A and B when placed at certain distance in air repel each other with a force of  $F$ . Another identical uncharged sphere C is first placed in contact with A and then in contact with B and finally placed at midpoint between spheres A and B. The force experienced by sphere C will be:

A  $3F/2$

B  $3F/4$

C  $F$

D  $2F$


NTA Ans. (B)

Reso Ans. (B)

Sol. Initially  $F = \frac{kq^2}{r^2}$  q- charge on A & B.

When A & C are touched. Charge on each becomes  $\frac{q}{2}$

Now when B & C are touched each becomes  $\frac{3q}{4}$



$$F_C = F_{AC} - F_{BC} = \frac{k \frac{q}{2} \frac{3q}{4}}{\left(\frac{r}{2}\right)^2} - \frac{k \left(\frac{3q}{4}\right)^2}{\left(\frac{r}{2}\right)^2}$$

$$= \frac{ka^2}{r^2} \left( \frac{3}{2} - \frac{9}{4} \right) = \frac{ka^2}{r^2} \left( -\frac{3}{4} \right)$$

$$(F_C) = \frac{3F}{4}$$

**Resonance Eduventures Ltd.**

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | 7340010333 | [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) | [twitter.com/ResonanceEdu](https://twitter.com/ResonanceEdu) | [www.youtube.com/resowatch](https://www.youtube.com/resowatch) | [blog.resonance.ac.in](https://blog.resonance.ac.in)

2. Match List I with List II.

List I	List II
A. Torque	I. $\text{Nms}^{-1}$
B. Stress	II. $\text{J kg}^{-1}$
C. Latent Heat	III. $\text{Nm}$
D. Power	IV. $\text{Nm}^{-2}$

Choose the correct answer from the options given below:

- A A-III, B-II, C-I, D-IV
- B A-III, B-IV, C-II, D-I
- C A-IV, B-I, C-III, D-II
- D A-II, B-III, C-I, D-IV

NTA Ans. (B)

Reso Ans. (B)

3. Two identical thin metal plates has charge  $q_1$  and  $q_2$  respectively such that  $q_1 > q_2$ . The plates were brought close to each other to form a parallel plate capacitor of capacitance  $C$ . The potential difference between them is :

- A  $\frac{(q_1 + q_2)}{C}$
- B  $\frac{(q_1 - q_2)}{C}$
- C  $\frac{(q_1 - q_2)}{2C}$
- D  $\frac{2(q_1 - q_2)}{C}$

NTA Ans. (C)






Reso Ans. (C)

## Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555  7340010333  facebook.com/ResonanceEdu  twitter.com/ResonanceEdu  www.youtube.com/resowatch  blog.resonance.ac.in

Sol.

$$V = \frac{q}{C} = \frac{q_1 - q_2}{2C}$$

4. Given below are two statements: one is labelled as **Assertion A** and the other is labelled as **Reason R**.

**Assertion A:** Alloys such as constantan and manganin are used in making standard resistance coils.

**Reason R:** Constantan and manganin have very small value of temperature coefficient of resistance.

In the light of the above statements, choose the *correct* answer from the options

- A Both A and R are true and R is the correct explanation of A.
- B Both A and R are true but R is NOT the correct explanation of A.
- C A is true but R is false.
- D A is false but R is true.

NTA Ans. (A)

Reso Ans. (A)

Sol. Theory Based

5. A 1 m long wire is broken into two unequal parts X and Y. The X part of the wire is stretched into another wire W. Length of W is twice the length of X and the resistance of W is twice that of Y. Find the ratio of length of X and Y.

- A 1:4
- B 1:2
- C 4:1
- D 2:1

## Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | 7340010333 | [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) | [twitter.com/ResonanceEdu](https://twitter.com/ResonanceEdu) | [www.youtube.com/resowatch](https://www.youtube.com/resowatch) | [blog.resonance.ac.in](https://blog.resonance.ac.in)



NTA Ans. (B)

Reso Ans. (B)

Sol. Now length of x is doubled its area becomes half.


$$R_x^1 = \int \frac{2l}{A_1} = 4 \int \frac{l}{A}$$

Given  $R_x^1 = 2R_y$


$$4 \int \frac{l}{A} = 2 \int \frac{(1-l)}{A}$$

$$\frac{l}{1-l} = \frac{1}{2}$$

$$\frac{x}{y} = \frac{1}{2}$$



6. A wire X of length 50 cm carrying a current of 2 A is placed parallel to a long wire Y of length 5 m. The wire Y carries a current of 3 A. The distance between two wires is 5 cm and currents flow in the same direction. The force acting on the wire Y is



A  $1.2 \times 10^{-5}$  N directed towards wire X.

B  $1.2 \times 10^{-4}$  N directed away from wire X.

C  $1.2 \times 10^{-4}$  N directed towards wire X.

D  $2.4 \times 10^{-5}$  N directed towards wire X.

NTA Ans. (A)

Reso Ans. (A)

## Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 7340010333 facebook.com/ResonanceEdu twitter.com/ResonanceEdu www.youtube.com/resowatch blog.resonance.ac.in

Sol.  $F = I_1 l B_2$

$$= 2(0.5) \left[ \frac{\mu_0(I_2)}{2\pi d} \right]$$

$$= 2(0.5) \left[ \frac{2 \times 10^{-7} \times 3}{5 \times 10^{-2}} \right]$$

$$= \frac{6}{5} \times 10^{-5} \text{ N}$$

$$= 1.2 \times 10^{-5} \text{ N, attraction}$$

7. A juggler throws balls vertically upwards with same initial velocity in air. When the first ball reaches its highest position, he throws the next ball. Assuming the juggler throws  $n$  balls per second, the maximum height the balls can reach is

A  $g/2n$

B  $g/n$

C  $2gn$

D  $g/2n^2$

NTA Ans. (D)

Reso Ans. (D)

Sol. Time taken by each ball to reach maximum height



$n$  ball per/sec

$$\frac{u}{g} = \frac{1}{n}$$

Maximum height of each ball

$$H = \frac{u^2}{2g} = \frac{g^2}{2n^2 \times g}$$






$$= \frac{g}{2n^2}$$

## Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555  7340010333  facebook.com/ResonanceEdu  twitter.com/ResonanceEdu  www.youtube.com/resowatch  blog.resonance.ac.in

8. A circuit element X when connected to an a.c. supply of peak voltage 100 V gives a peak current of 5 A which is in phase with the voltage. A second element Y when connected to the same a.c. supply also gives the same value of peak current which lags behind the voltage by  $\frac{\pi}{2}$ . If X and Y are connected in series to the same supply, what will be the rms value of the current in ampere?

A  $\frac{10}{\sqrt{2}}$


B  $\frac{5}{\sqrt{2}}$

C  $5\sqrt{2}$

D  $\frac{5}{2}$

NTA Ans. (D)

Reso Ans. (D)



$$V_{\max} = 100V$$

$$R = \frac{V}{I} = \frac{100}{5} = 20\Omega$$


$$I_{\max} = 5A$$

y with  $90^\circ$  phase

$$V_{\max} = 100V$$

$$x = \frac{V}{I} = \frac{100}{5} = 20\Omega$$

when X and Y are connected in series with source



$$V_{\max} = 100$$

$$i_{\max} = \frac{V_{\max}}{Z} = \frac{100}{\sqrt{x^2 + y^2}} = \frac{100}{\sqrt{20^2 + 20^2}}$$

$$i_{\max} = \frac{5}{\sqrt{2}}, \quad i_{\text{rms}} = \frac{5}{2}$$

## Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | 7340010333 | [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) | [twitter.com/ResonanceEdu](https://twitter.com/ResonanceEdu) | [www.youtube.com/resowatch](https://www.youtube.com/resowatch) | [blog.resonance.ac.in](https://blog.resonance.ac.in)

9. An unpolarised light beam of intensity  $2I_0$  is passed through a polaroid P and then through another polaroid Q which is oriented in such a way that its passing axis makes an angle of  $30^\circ$  relative to that of P. The intensity of the emergent light is

A  $\frac{I_0}{4}$

B  $\frac{I_0}{2}$


C  $\frac{3I_0}{4}$

D  $\frac{3I_0}{2}$

NTA Ans. (C)

Reso Ans. (C)

Sol.



10. An  $\alpha$  particle and a proton are accelerated from rest through the same potential difference. The ratio of linear momenta acquired by above two particles will be:

A  $\sqrt{2}:1$

B  $2\sqrt{2}:1$

C  $4\sqrt{2}:1$

D  $8:1$

NTA Ans. (B)

Reso Ans. (B)

## Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 7340010333 facebook.com/ResonanceEdu twitter.com/ResonanceEdu www.youtube.com/resowatch blog.resonance.ac.in



**Sol.** Momentum of acceleration charged particle.

$$p = \sqrt{2meV}$$

$$p_p = \sqrt{2meV} \quad p_\alpha = \sqrt{2 \times (4m)(2e)V}$$

$$\frac{p_\alpha}{p_p} = \frac{\sqrt{8}}{1} = 2\sqrt{2} : 1$$

11. Read the following statements:

- (A) Volume of the nucleus is directly proportional to the mass number.
- (B) Volume of the nucleus is independent of mass number.
- (C) Density of the nucleus is directly proportional to the mass number.
- (D) Density of the nucleus is directly proportional to the cube root of the mass number.
- (E) Density of the nucleus is independent of the mass number.

Choose the correct option from the following options.

- A (A) and (D) only.
- B (A) and (E) only.
- C (B) and (E) only.
- D (A) and (C) only.

NTA Ans. (B)

Reso Ans. (B)

**Sol.**  $r = r_0 A^{1/3}$

$$r^3 \propto A$$

$$\rho = M/V = \frac{Am_p}{A \times \frac{4}{3} \pi r_0^3} = \frac{m_p}{\frac{4}{3} \pi r_0^3}$$

12. An object of mass 1 kg is taken to a height from the surface of earth which is equal to three times the radius of earth. The gain in potential energy of the object will be [If,  $g=10\text{ms}^{-2}$  and radius of earth = 6400 km]

- A 48 MJ
- B 24 MJ
- C 36 MJ
- D 12 MJ

NTA Ans. (A)






Reso Ans. (A)

## Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555  7340010333  facebook.com/ResonanceEdu  twitter.com/ResonanceEdu  www.youtube.com/resowatch  blog.resonance.ac.in

Sol.  $u_i = m \left[ -\frac{GM}{R} \right]$

$$u_f = m \left[ -\frac{GM}{4R} \right]$$

$$\Delta u = u_f - u_i$$

$$\Delta u = \frac{GMm}{R} - \frac{GMm}{4R}$$

$$\Delta u = \frac{3}{4} \frac{GMm}{R}$$

13. A ball is released from a height  $h$ . If  $t_1$  and  $t_2$  be the time required to complete first half and second half of the distance respectively. Then, choose the correct relation between  $t_1$  and  $t_2$ .

A  $t_1 = (\sqrt{2})t_2$

B  $t_1 = (\sqrt{2} - 1)t_2$


C  $t_2 = (\sqrt{2} + 1)t_1$

D  $t_2 = (\sqrt{2} - 1)t_1$

NTA Ans. (D)

Reso Ans. (D)

Sol.



$$\frac{h}{2} = \frac{1}{2} g t_1^2$$

$$h = \frac{1}{2} g (t_1 + t_2)^2$$

$$2 \frac{1}{2} g t_1^2 = \frac{1}{2} g (t_1 + t_2)^2$$

$$\sqrt{2} t_1 = t_1 + t_2 \quad (\sqrt{2} - 1) t_1 = t_2$$

## Resonance Eduventures Ltd.


Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 7340010333 facebook.com/ResonanceEdu twitter.com/ResonanceEdu www.youtube.com/resowatch blog.resonance.ac.in

14. Two bodies of masses  $m_1 = 5 \text{ kg}$  and  $m_2 = 3 \text{ kg}$  are connected by a light string going over a smooth light pulley on a smooth inclined plane as shown in the figure. The system is at rest. The force exerted by the inclined plane on the body of mass  $m_1$  will be : [Take  $g = 10 \text{ ms}^{-2}$ ]




- A 30 N  
B 40 N  
C 50 N  
D 60 N

NTA Ans. (B)

Reso Ans.

Sol.  $T = 30 \text{ N}$

$$T = 50 \sin \theta$$



$$30 = 50 \sin \theta \Rightarrow \sin \theta = \frac{3}{5} \quad \theta = 37^\circ$$

$$N = 50 \cos 37^\circ = 40 \text{ N}$$

15. If momentum of a body is increased by 20%, then its kinetic energy increases by

- A 36%  
B 40%  
C 44%  
D 48%

NTA Ans. (C)

Reso Ans. (C)

## Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | 7340010333 | [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) | [twitter.com/ResonanceEdu](https://twitter.com/ResonanceEdu) | [www.youtube.com/resowatch](https://www.youtube.com/resowatch) | [blog.resonance.ac.in](https://blog.resonance.ac.in)

Sol. Initially  $K_i = \frac{p^2}{2m}$

Momentum is increase by 20%

$$p_f = p + 0.20p = 1.20p$$

$$K_f = \frac{(1.20p)^2}{2m} = \frac{1.44p^2}{2m}$$

$$\frac{\Delta KE}{K_i} = \frac{\frac{1.44p^2}{2m} - \frac{p^2}{2m}}{\frac{p^2}{2m}} = 0.44$$

$$\frac{\Delta KE}{K_i} \times 100 = 44\%$$

16. The torque of a force  $5\hat{i} + 3\hat{j} - 7\hat{k}$  about the origin is  $\tau$ . If the force acts on a particle whose position vector is  $2\hat{i} + 2\hat{j} + \hat{k}$ , then the value of  $\tau$  will be

A  $11\hat{i} + 19\hat{j} - 4\hat{k}$

B  $-11\hat{i} + 9\hat{j} - 16\hat{k}$

C  $-17\hat{i} + 19\hat{j} - 4\hat{k}$

D  $17\hat{i} + 9\hat{j} + 16\hat{k}$

NTA Ans. (C)

Reso Ans. (C)






Sol.  $\vec{\tau} = \vec{r} \times \vec{f} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 2 & 1 \\ 5 & 3 & -7 \end{vmatrix} = -17\hat{i} + 19\hat{j} - 4\hat{k}$

## Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005


Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555  7340010333  facebook.com/ResonanceEdu  twitter.com/ResonanceEdu  www.youtube.com/resowatch  blog.resonance.ac.in



17. A thermodynamic system is taken from an original state D to an intermediate state E by the linear process shown in the figure. Its volume is then reduced to the original volume from E to F by an isobaric process. The total work done by the gas from D to E to F will be



- A -450 J  
B 450 J  
C 900 J  
D 1350 J

NTA Ans. (B)

Reso Ans. (B)

Sol.  $\Delta w = \text{area enclosed in the cycle}$   
 $= \frac{1}{2} \times 3 \times 300 = 450 \text{ J}$

18. The vertical component of the earth's magnetic field is  $6 \times 10^{-5} \text{ T}$  at any place where the angle of dip is  $37^\circ$ . The earth's resultant magnetic field at that place will be (Given  $\tan 37^\circ = \frac{3}{4}$ )

- A  $8 \times 10^{-5} \text{ T}$   
B  $6 \times 10^{-5} \text{ T}$   
C  $5 \times 10^{-4} \text{ T}$   
D  $1 \times 10^{-4} \text{ T}$

NTA Ans. (D)

Reso Ans. (D)

Sol.  $\tan \lambda = \frac{B_V}{B_H}$   
 $\tan 37^\circ = \frac{6 \times 10^{-5}}{B_H} = B_H = 8 \times 10^{-5} \text{ T}$   
 $B_{\text{Net}} = \sqrt{B_H^2 + B_V^2} = 10 \times 10^{-5} = 10^{-4} \text{ T}$

## Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 7340010333 facebook.com/ResonanceEdu twitter.com/ResonanceEdu www.youtube.com/resowatch blog.resonance.ac.in

19. The root mean square speed of smoke particles of mass  $5 \times 10^{-17}$  kg in their Brownian motion in air at NTP is approximately. [Given  $k = 1.38 \times 10^{-23}$  JK<sup>-1</sup>]

- A  $60 \text{ mm s}^{-1}$
- B  $12 \text{ mm s}^{-1}$
- C  $15 \text{ mm s}^{-1}$
- D  $36 \text{ mm s}^{-1}$

NTA Ans. (C)

Reso Ans. (C)

Sol.  $V_{\text{rms}} = \sqrt{\frac{3kT}{m}} \approx 15 \text{ mm/s}$

20. Light enters from air into a given medium at an angle of  $45^\circ$  with interface of the air-medium surface. After refraction, the light ray is deviated through an angle of  $15^\circ$  from its original direction. The refractive index of the medium is :


- A 1.732
- B 1.333
- C 1.414
- D 2.732

NTA Ans. (C)

Reso Ans. (C)

Sol.  $i = 45^\circ$   
 $\Delta = 15^\circ$   
 $\therefore r = 30^\circ$

$$\mu = \frac{\sin i}{\sin r} = \frac{\sin 45}{\sin 30} = \frac{\frac{1}{\sqrt{2}}}{\frac{1}{2}} = \sqrt{2} = 1.414$$



## Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222


To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 7340010333 facebook.com/ResonanceEdu twitter.com/ResonanceEdu www.youtube.com/resowatch blog.resonance.ac.in

21. A tube of length 50 cm is filled completely with an incompressible liquid of mass 250 g and closed at both ends. The tube is then rotated in horizontal plane about one of its ends with a uniform angular velocity  $x\sqrt{F}$  rad s<sup>-1</sup>. If F be the force exerted by the liquid at the other end then the value of x will be \_\_\_\_\_.

NTA Ans. 4

Reso Ans. 4



$$F = \int (dm)\omega^2 x$$

$$= \int_0^L \left(\frac{m}{L} dx\right) \omega^2 x$$

$$= \frac{m}{L} \omega^2 \frac{L^2}{2}$$

$$= \frac{m\omega^2 L}{2}$$

$$\omega = \sqrt{\frac{2}{mL} F}$$

$$= \sqrt{\frac{2}{0.25 \times 0.5}} \sqrt{F}$$

$$= \sqrt{16} \sqrt{F}$$

$$= 4\sqrt{F}$$

22. Nearly 10% of the power of a 110 W light bulb is converted to visible radiation. The change in average intensities of visible radiation, at a distance of 1 m from the bulb to a distance of 5 m is  $a \times 10^{-2}$  W/m<sup>2</sup>. The value of 'a' will be \_\_\_\_\_.

NTA Ans. 84

Reso Ans. 84

Sol.  $P' = 10\%$  of 110 W

$$= \frac{10}{100} \times 110 \text{ W} = 11 \text{ W}$$

$$I_1 - I_2 = \frac{P'}{4\pi r_1^2} - \frac{P}{4\pi r_2^2} = \frac{11}{4\pi} \left[ \frac{1}{1} - \frac{1}{25} \right]$$

$$= \frac{11}{4\pi} \times \frac{24}{25} = \frac{264}{\pi} \times 10^{-2} = 84 \times 10^{-2} \text{ W/m}^2$$

## Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | WhatsApp : 7340010333 | Facebook : facebook.com/ResonanceEdu | Twitter : twitter.com/ResonanceEdu | YouTube : www.youtube.com/resowatch | Blog : blog.resonance.ac.in



23. A metal wire of length 0.5 m and cross-sectional area  $10^{-4} \text{ m}^2$  has breaking stress  $5 \times 10^8 \text{ Nm}^{-2}$ . A block of 10 kg is attached at one end of the string and is rotating in a horizontal circle. The maximum linear velocity of block will be \_\_\_  $\text{ms}^{-1}$ .

NTA Ans. 50

Reso Ans. 50

Sol.  $T = \frac{mv^2}{\ell} = \frac{10 \times v^2}{0.5} = 20v^2$

$T_{\max} = \text{Breaking stress} \times \text{Area}$

$= 5 \times 10^8 \times 10^{-4} = 5 \times 10^4$

$20v^2 = 5 \times 10^4$

$V = \sqrt{\frac{1}{4} \times 10^4} = 50 \text{ m/s}$

24. The velocity of a small ball of mass 0.3 g and density 8 g/cc when dropped in a container filled with glycerine becomes constant after some time. If the density of glycerine is 1.3 g/cc, then the value of viscous force acting on the ball will be  $x \times 10^{-4} \text{ N}$ , The value of x is \_\_\_\_\_. [use  $g = 10 \text{ m/s}^2$ ]

NTA Ans. 25

Reso Ans. 25

Sol.  $mg - F_b - F_v = 0$

$F_v = mg - F_b$

$F_v = \frac{4}{3} \pi r^3 \rho_b g - \frac{4}{3} \pi r^3 \rho_g g$

$= \rho_b \frac{4}{3} \pi r^3 g \left( 1 - \frac{\rho_g}{\rho_b} \right)$

$= mg \left( 1 - \frac{\rho_g}{\rho_b} \right)$

$= 0.3 \times 10^{-3} \times 10 \left( 1 - \frac{1.3}{8} \right)$

$= 2.5 \times 10^{-3} \text{ N}$

25. A modulating signal  $2\sin(6.28 \times 10^6)t$  is added to the carrier signal  $4\sin(12.56 \times 10^9)t$  for amplitude modulation. The combined signal is passed through a non-linear square law device. The output is then passed through a band pass filter. The bandwidth of the output signal of band pass filter will be \_\_\_ MHz.

NTA Ans. 2






Reso Ans. 2

## Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555  7340010333  facebook.com/ResonanceEdu  twitter.com/ResonanceEdu  www.youtube.com/resowatch  blog.resonance.ac.in



**Sol.** Frequencies present

$$2f_c, f_c + f_m, f_c, f_c - f_m, 2f_m, f_m$$

After passing through band pass filter.

$$f_c + f_m, f_c, f_c - f_m$$

$$\text{Band width} = 2f_m = \frac{2\omega_m}{2\pi} = 2 \text{ MHz}$$

26. The speed of a transverse wave passing through a string of length 50 cm and mass 10 g is  $60 \text{ ms}^{-1}$ . The area of cross-section of the wire is  $2.0 \text{ mm}^2$  and its Young's modulus is  $1.2 \times 10^{11} \text{ Nm}^{-2}$ . The extension of the wire over its natural length due to its tension will be  $x \times 10^{-5} \text{ m}$ . The value of  $x$  is     .

**NTA Ans.** 15

**Reso Ans.** 15

**Sol.**

$$V_\omega = \sqrt{\frac{T}{\mu}}$$

$$60 = \sqrt{\frac{T}{10 \times 10^{-3}}} \times 0.5$$

$$T = \frac{(60)^2 \times 10^{-2}}{0.5} = 72 \text{ N}$$

$$\Delta l = \frac{F l}{A y} = \frac{72 \times 0.5}{2 \times 10^{-6} \times 1.2 \times 10^{11}}$$

$$= \frac{72 \times 5}{24} \times 10^{-5} = 15 \times 10^{-5}$$

27. The metallic bob of simple pendulum has the relative density 5. The time period of this pendulum is 10 s. If the metallic bob is immersed in water, then the new time period becomes  $5\sqrt{x}$  s. The value of  $x$  will be     .

**NTA Ans.** 5

**Reso Ans.** 5

**Sol.**

$$\frac{T_f}{T} = \sqrt{\frac{g}{g\left(1 - \frac{\rho_\ell}{\rho}\right)}} = \sqrt{\frac{1}{1 - \frac{1}{5}}} = \frac{\sqrt{5}}{2}$$

$$T_f = 5\sqrt{5}$$

$$x = 5$$

## Resonance Eduventures Ltd.


Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | 7340010333 | [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) | [twitter.com/ResonanceEdu](https://twitter.com/ResonanceEdu) | [www.youtube.com/resowatch](https://www.youtube.com/resowatch) | [blog.resonance.ac.in](https://blog.resonance.ac.in)


28. A 8 V Zener diode along with a series resistance R is connected across a 20 V supply (as shown in the figure). If the maximum Zener current is 25 mA, then the minimum value of R will be \_\_\_\_\_  $\Omega$ .



NTA Ans. 480

Reso Ans. 480

Sol.



$$i = \frac{V}{R_{\min}}$$

$$R_{\min} = \frac{V}{i} = \frac{12}{25 \times 10^{-3}} = 480 \Omega$$

29. Two radioactive materials A and B have decay constants  $25\lambda$  and  $16\lambda$  respectively. If initially they have the same number of nuclei, then the ratio of the number of nuclei of B to that of A will be "e" after a time  $\frac{1}{a\lambda}$ . The value of a is \_\_\_\_\_.

NTA Ans. 9

Reso Ans. 9

Sol.  $N_0 \xrightarrow{16\lambda} N_1 = N_0 e^{-16\lambda t}$   
 $t = 0 \qquad t = t$

$N_0 \xrightarrow{25\lambda} N_2 = N_0 e^{-25\lambda t}$   
 $t = 0 \qquad t = t$

Given  $\frac{N_1}{N_2} = e$

$$\frac{N_0 e^{-16\lambda t}}{N_0 e^{-25\lambda t}} = e$$

$$\Rightarrow e^{9\lambda t} = e$$

$$\Rightarrow 9\lambda t = 1$$

$$\Rightarrow t = \frac{1}{9\lambda}$$

Ans. a = 9

## Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555 | 7340010333 | [facebook.com/ResonanceEdu](https://www.facebook.com/ResonanceEdu) | [twitter.com/ResonanceEdu](https://twitter.com/ResonanceEdu) | [www.youtube.com/resowatch](https://www.youtube.com/resowatch) | [blog.resonance.ac.in](http://blog.resonance.ac.in)

30. A capacitor of capacitance  $500 \mu\text{F}$  is charged completely using a dc supply of  $100 \text{ V}$ . It is now connected to an inductor of inductance  $50 \text{ mH}$  to form an LC circuit. The maximum current in LC circuit will be \_\_\_ A.

NTA Ans. 10

Reso Ans. 10

Sol.  $q = 500 \times 10^{-6} \times 100 = 5 \times 10^{-2} \text{ C}$

If connected to LC

$$\frac{q^2}{2C} = \frac{1}{2} Li^2$$






$$i = \frac{q}{\sqrt{LC}} = \frac{5 \times 10^{-2}}{\sqrt{50 \times 10^{-3} \times 500 \times 10^{-6}}} = \frac{5 \times 10^{-2}}{5 \times 10^{-3}} = 10 \text{ A}$$

## Resonance Eduventures Ltd.

Reg. Office & Corp. Office : CG Tower, A-46 & 52, IPIA, Near City Mall, Jhalawar Road, Kota (Raj.) - 324005

Ph. No.: +91-744-2777777, 2777700 | FAX No. : +91-022-39167222

To Know more : sms RESO at 56677 | Website : www.resonance.ac.in | E-mail : contact@resonance.ac.in | CIN : U80302RJ2007PLC024029

Toll Free : 1800 258 5555  7340010333  facebook.com/ResonanceEdu  twitter.com/ResonanceEdu  www.youtube.com/resowatch  blog.resonance.ac.in



Numbers that Inspire Students to EXCEL



\*Since 2001 | \*\*Students Qualified from JEE (Main) to JEE (Advanced) since 2012 | Total Selections in AIR in TOP-100 mentioned are in JEE (Adv.) / JEE (Main) since 2002, JEE (Main) / AIEEE (since 2005), NEET (UG) / APJKT (since 2012) | AIR: All India Rank


The 10-Building Blocks of Proven & Trusted Teaching Methodology @ Resonance

Success is Assured  
when you Do\*

200 X Practice | 20 X Testing

<b>1500+</b> Hrs. Classroom Teaching	<b>10000+</b> Pages Study Material
<b>45000+</b> Questions Practice Problems	<b>50+</b> Tests   <b>200+</b> Hrs. Testing & Assessment

\* As per Latest Pattern of JEE (Main) & JEE (Advanced)



The figures (approx.) shown in the graph are of 2 Years Classroom Program (VIKAAS-XI & VIETA-XII) for JEE (Advanced) @ Resonance in Academic Session 2021-22. The figures vary for JEE (Main), NEET (UG) and Other Courses.

The Strong Faculty Team at Resonance Kota to deliver this successful Teaching Methodology



Photo Taken on 20<sup>th</sup> June 2022 | Some Faculty Members were not present in the Photo Session.

SCHOLARSHIP UPTO

**100%**

Based on JEE (Main) 2022 NTA Score (Percentile) & Scholarship Test (ResoNET)

Admission Announcement: 2022-23 | Class: 5 to 12 & 12+

**ResoNET** 3<sup>rd</sup> & 10<sup>th</sup> July

Target: JEE (Advanced) | JEE (Main) | NEET (UG) | Pre-Foundation (V to X) | Board

Polish your subject knowledge to Shine in JEE (Advanced) 2022 with the guidance of HODs & Top Notch Sr. Faculty of Resonance

**SPARK** 7 WEEKS COMPACT COURSE OFFLINE/ONLINE

from 4<sup>th</sup> July 2022

Scholarship upto 90%